

# THE RAMSEY-TYPE VERSION OF A PROBLEM OF POMERANCE AND SCHINZEL

PÉTER PÁL PACH

*The author would like to dedicate this paper to Andrzej Schinzel, on the occasion of his upcoming 75th birthday.*

ABSTRACT. We prove that for any  $r$ -colouring of the squarefree numbers the equation  $a_1 a_2 \dots a_k = b_1 b_2 \dots b_l$  has a primitive monochromatic solution.

## 1. INTRODUCTION

A set  $H$  is called product free if  $a, b \in H$  implies  $ab \notin H$ . Hajdu, Schinzel and Skalba have shown that a product free subset of the positive integers can have upper density arbitrarily close to 1 [4]. Sárközy has suggested to investigate the Ramsey-type variation of the problem: is it true that for any  $r$ -colouring of  $\mathbb{N}$  the equation  $ab = c$  has a monochromatic solution different from the trivial solution  $1 \cdot 1 = 1$ . In particular he asked the question for squarefree numbers:

**Problem 1.** *Is it true that for any  $r$ -colouring of the squarefree numbers greater than 1 the equation  $ab = c$  has a monochromatic solution?*

There are several other questions without density theorems, where the Ramsey-type version was answered positively, see for example [1], [5]. It is a consequence of Schur's theorem [9] that Sárközy's original problem always has a solution among the powers of 2.

**Proposition 1.** *For every  $r$ -colouring of the 2-powers the equation  $ab = c$  has a nontrivial solution.*

*Proof.* Let us colour the 2-powers by  $r$  colours. We define a colouring of  $\mathbb{N}$  by  $r$  colours in the following way. Let the colour of  $x \in \mathbb{N}$  be the colour of  $2^x$ . By Schur's theorem the equation  $x + y = z$  has a monochromatic solution in  $\mathbb{N}$ . Then the equation  $ab = c$  also has a monochromatic solution (for the original colouring) among the 2-powers, namely  $a = 2^x, b = 2^y, c = 2^z$ .  $\square$

Pomerance and Schinzel has proved that for Problem 1 the answer is affirmative if  $r = 2$  ([7]). In this paper we settle the problem for arbitrary  $r$ , and extend the results for more general equations. We show that the equation  $a_1 a_2 \dots a_k = b_1 b_2 \dots b_l$  has a nontrivial monochromatic solution for every  $r$ -colouring of the squarefree numbers.

---

2010 *Mathematics Subject Classification*: Primary 11B75.

*Key words and phrases*: algebraic equation, Ramsey type, solution set.

## 2. SQUAREFREE NUMBERS

The result of Hajdu, Schinzel and Skałba implies that there is no density theorem for the equation  $ab = c$ . The following example shows that if  $k \neq l$ , then there is no density theorem for the equation  $a_1 a_2 \dots a_k = b_1 b_2 \dots b_l$ , either.

**Example 2.1.** Let  $A_n = \{4i+2 : 0 \leq i, 4i+2 \leq n\}$ . If  $a_1, a_2, \dots, a_k, b_1, b_2, \dots, b_l \in A_n$ , then the exponent of 2 is  $k$  in the canonical form of  $a_1 a_2 \dots a_k$  and  $l$  in  $b_1 b_2 \dots b_l$ . Thus the equation  $a_1 a_2 \dots a_k = b_1 b_2 \dots b_l$  doesn't have a solution in  $A_n$  if  $k \neq l$ . The size of  $A_n$  is  $\frac{1}{4} \cdot n + O(1)$ .

If  $k = l$ , then  $a_1 = \dots = a_k = b_1 = \dots = b_k$  is a solution. We say that  $a_1, a_2, \dots, a_k, b_1, b_2, \dots, b_l$  is a primitive solution of the equation  $a_1 a_2 \dots a_k = b_1 b_2 \dots b_l$  if  $a_1, a_2, \dots, a_k, b_1, b_2, \dots, b_l$  are pairwise distinct. From now we look only for primitive solutions of equations.

In case  $k = l$  there is a density theorem for primitive solutions if  $k$  and  $l$  are even.

**Proposition 2.** *Let  $k \in \mathbb{N}$  be even. For arbitrary  $\varepsilon > 0$  there exists  $N = N(\varepsilon)$  such that for every  $n \geq N$  and  $A \subseteq \{1, \dots, n\}$  with size  $|A| \geq \varepsilon n$  the equation  $a_1 a_2 \dots a_k = b_1 b_2 \dots b_k$  has a primitive solution in  $A$ .*

*Proof.* The proof is by induction on  $k$ . At first let  $k = 2$  and  $\varepsilon > 0$  be arbitrary. The bound  $N$  is chosen later. Let  $A \subseteq \{1, \dots, n\}$ , where  $n \geq N$  and  $|A| > \varepsilon n$ . In [2] it is proved that only  $o(n^2)$  numbers can be found in the "multiplication table" of the integers up to  $n$ . As  $A \subseteq \{1, \dots, n\}$ , the set  $A \cdot A = \{c_1 c_2 : c_1, c_2 \in A\}$  has at most  $o(n^2)$  elements. There are  $\binom{|A|}{2} = \frac{\varepsilon^2}{2} \cdot n^2 + o(n^2)$  pairs  $c_1, c_2$  with  $c_1, c_2 \in A$  and  $c_1 \neq c_2$ . Now, choose  $N$  such that  $\binom{|A|}{2}$  is larger than the size of  $A \cdot A$ . Thus there exists an element in  $A \cdot A$  which can be written as a product of two different elements of  $A$  in at least two different ways:  $a_1 a_2 = b_1 b_2$ . This way we obtained a primitive solution.

Now, assume that  $4 \leq k \in 2\mathbb{N}$  and the statement holds for  $k - 2$ . Let  $\varepsilon > 0$  be arbitrary. By the induction hypothesis there exists some  $N$  such that for any set  $B \subseteq \{1, \dots, n\}$  with at least  $\frac{\varepsilon}{3} \cdot n$  elements, the equations  $a_1 a_2 \dots a_{k-2} = b_1 b_2 \dots b_{k-2}$  and  $a_{k-1} a_k = b_{k-1} b_k$  have a primitive solution in  $B$  if  $n \geq N$ . Let  $A \subseteq \{1, \dots, n\}$  having at least  $\varepsilon n$  elements. If  $n \geq 3/\varepsilon$ , then  $A$  can be partitioned into two disjoint parts  $A_1$  and  $A_2$  both of size at least  $\frac{\varepsilon}{3} \cdot n$ . If  $n \geq N$ , then  $a_1 a_2 \dots a_{k-2} = b_1 b_2 \dots b_{k-2}$  has a primitive solution in  $A_1$  and  $a_{k-1} a_k = b_{k-1} b_k$  has a primitive solution in  $A_2$ . Therefore,  $a_1, a_2, \dots, a_k, b_1, b_2, \dots, b_k$  is a primitive solution of  $a_1, a_2, \dots, a_k = b_1, b_2, \dots, b_k$  in  $A$ . □

The case when  $k = l$  is odd is still open.

**Problem 2.** *Is it true that for every odd  $k > 1$  and  $\varepsilon > 0$  there exists some  $N$  such that for every  $N \leq n$  and  $A \subseteq \{1, 2, \dots, n\}$  with size at least  $\varepsilon n$  the equation  $a_1 a_2 \dots a_k = b_1 b_2 \dots b_k$  has a primitive solution in  $A$ ?*

For the main result of the paper the following form of Ramsey's theorem will be used ([3], [6]):

**Ramsey's Theorem.** *Let  $r$  and  $t$  be positive integers. Let us colour the at most  $t$ -element subsets of a set  $S$  by  $r$  colours. Then for every positive integer  $n$  there exists a positive integer  $d$  such that if  $|S| > d$ , then  $S$  has a subset  $H$  with  $n$  elements, such that any two subsets of the same size not greater than  $t$  have the same colour, that is, for every  $H_1, H_2 \subseteq H$ ,  $|H_1| = |H_2| \leq t$  the colour of  $H_1$  and  $H_2$  are the same.*

By Ramsey's theorem for every  $n$  there exists  $d$  such that if  $|S| > d$ , then there exists a subset  $H \subseteq S$ ,  $|H| = n$  such that every one-element subset of  $H$  has the same colour, every two-element subset of  $H$  has the same colour, and so on, every subset of  $H$  with  $t$  elements has the same colour. The bound for this integer  $d$  is called a Ramsey-number and the best known bound is multiply exponential in  $r$ .

The following version of Rado's theorem is also needed ([6],[8]):

**Rado's Theorem.** *Let  $v \geq 2$ . Let  $c_i \in \mathbb{Z} \setminus \{0\}$ ,  $1 \leq i \leq v$  be constants such that there exists a nonempty  $D \subseteq \{c_i : 1 \leq i \leq v\}$  such that  $\sum_{d \in D} d = 0$ . If there exist distinct integers (not necessarily positive)  $y_i$  such that  $\sum c_i y_i = 0$ , then for every natural number  $r$  there exists some  $t$  such that for every  $r$ -colouring of the set  $\{1, 2, \dots, t\}$  the equation*

$$c_1 x_1 + \dots + c_v x_v = 0$$

*has a monochromatic solution  $b_1, \dots, b_v$  in  $\{1, 2, \dots, t\}$ , where the  $b_i$ -s are distinct.*

Now we prove that for every  $r$ -colouring of the squarefree numbers the equation  $a_1 a_2 \dots a_k = b_1 b_2 \dots b_l$  has a primitive monochromatic solution if  $k \geq 2$ .

**Theorem 3.** *For every  $k \geq 2, l, r \in \mathbb{N}$  and every  $r$ -colouring of the square-free numbers greater than 1 the equation*

$$(1) \quad a_1 a_2 \dots a_k = b_1 b_2 \dots b_l$$

*has a primitive monochromatic solution.*

*Proof.* The squarefree numbers are in a one-to-one correspondence with the finite subsets of primes. To each squarefree number we assign the set of its prime divisors. The product of two squarefree numbers is squarefree if and only if the two sets are disjoint. Moreover, in this case the product corresponds to the union of the two subsets.

For a given  $r$ -colouring of the squarefree numbers we define a colouring of the finite subsets of primes. Each subset is coloured by the colour of the product of its elements. If we find nonempty subsets of primes  $A_1, \dots, A_k, B_1, \dots, B_l$  such that

- (i)  $\cup A_i = \cup B_j$ ,
- (ii)  $A_1, \dots, A_k, B_1, \dots, B_l$  are pairwise distinct,

then  $a_i = \prod_{p \in A_i} p$  for  $1 \leq i \leq k$  and  $b_j = \prod_{p \in B_j} p$  for  $1 \leq j \leq l$  is a primitive monochromatic solution of (1). Now we show that the sets  $A_i, B_j$  with the above conditions exist with the additional condition:

- (iii) the sizes  $|A_1| = \alpha_1, \dots, |A_k| = \alpha_k, |B_1| = \beta_1, \dots, |B_l| = \beta_l$  are distinct.

The equation

$$(2) \quad \alpha_1 + \dots + \alpha_k = \beta_1 + \dots + \beta_l$$

is equivalent to

$$\alpha_1 + \dots + \alpha_k - \beta_1 - \dots - \beta_l = 0,$$

hence Rado's theorem applies with  $v = k + l$ , and  $c_i = 1, y_i = i$  if  $1 \leq i \leq k$  and  $c_i = -1, y_i = -i$  if  $k < i < v$  and  $c_v = -1, y_v = \frac{(v-1)v}{2}$ . Let  $t$  be chosen such that for every  $r$ -colouring of  $\{1, 2, \dots, t\}$  the equation (2) has a monochromatic solution. Now, apply Ramsey's Theorem for this  $t$  and  $n = t \max(k, l)$ . There is a number  $d$  such that for every  $r$ -colouring of the subsets of the first  $d$  primes there is a subset of primes  $H$  such that  $|H| = n$ , and for every  $j \leq t$  the  $j$ -element subsets of  $H$  have the same colour. Let us colour the elements of the set  $\{1, \dots, t\}$  by  $r$  colours in the following way: for  $1 \leq i \leq t$  let the colour of  $i$  be the colour of the  $i$ -element subsets of  $H$ . By Rado's theorem there exist a monochromatic solution of (2). Let  $m = \alpha_1 + \dots + \alpha_k = \beta_1 + \dots + \beta_l$ , where  $\alpha_1, \dots, \alpha_k, \beta_1, \dots, \beta_l$  are distinct positive integers not greater than  $t$ . Consider an arbitrary partition  $A_1, \dots, A_k$  of type  $\alpha_1, \dots, \alpha_k$  and an arbitrary partition  $B_1, \dots, B_l$  of type  $\beta_1, \dots, \beta_l$  of the first  $m$  primes in  $H$ . These sets satisfy conditions (i)-(iii), so the statement is proved. □

### 3. ACKNOWLEDGEMENTS

The author is thankful to A. Sárközy for turning his attention to this topic and for the useful hints. The research of the author was supported by the Hungarian National Foundation for Scientific Research Grant K67870 and K100246.

### REFERENCES

- [1] P. Csikvári, K. Gyarmati, and A. Sárközy: *Density and Ramsey type results on algebraic equations with restricted solution sets* Integers, to appear
- [2] P. Erdős: *An asymptotic theorem in the theory of numbers*. Vestnik Leningrad Univ. 15 (1960), 41-49 (in Russian)
- [3] R. L. Graham, B. L. Rothschild, and J. H. Spencer. *Ramsey Theory*. Wiley-Interscience Series in Discrete Mathematics and Optimization. John Wiley & Sons, 1990.
- [4] L. Hajdu, A. Schinzel, M. Skalba: *Multiplicative properties of sets of positive integers*. Arch. Math. 93 (2009), 269-276.
- [5] N. Hindman: *Monochromatic sums equal to products in  $\mathbb{N}$* . Integers 11A (2011), Article 10, 1-10.
- [6] B. M. Landman and A. Robertson: *Ramsey Theory on the Integers*. American Mathematical Society, 2003.

- [7] C. Pomerance and A. Schinzel: *Multiplicative properties of sets of residues* Moscow Math. J. Combinatorics and Number Theory, to appear
- [8] R. Rado: *Studien zur Kombinatorik*. Math. Z. 36 (1) (1933), pp. 424-470.
- [9] I. Schur: *Über die Kongruenz  $x^m + y^m = z^m \pmod{p}$* . Jahresber Deutsch. Math. Verein. 25 (1916), 114-117. 5

EÖTVÖS LORÁND UNIVERSITY, DEPARTMENT OF ALGEBRA AND NUMBER THEORY,  
1117 BUDAPEST, PÁZMÁNY PÉTER SÉTÁNY 1/C, HUNGARY  
*E-mail address:* ppp24@cs.elte.hu